

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE	3. REPORT TYPE AND DATES COVERED Final, 30 Sep 88 to 29 Sep 90	
4. TITLE AND SUBTITLE High Resolution Spectroscopy of NF and BiF			5. FUNDING NUMBERS AFOSR-88-0347	
6. AUTHOR(S) Professor Roger Bacis				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Laboratoire de Spectrometrie Ionique et Moleculaire Universite Lyon I 43, Boulevard du 11 Novembre 1918 69622 Villeurbanne Cedex, France			8. PERFORMING ORGANIZATION REPORT NUMBER EOARD TR-91-04	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Sponsoring/Monitoring: European Office of Aerospace Research and Development, Box 14, FPO New York, NY 09510-0200			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; Distribution unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words)  During the period covered by the U.S. Air Force grant a Broida type oven was constructed allowing the spectroscopic investigation of the A O <sup>+</sup> and X O <sup>+</sup> electronic states of the BiF molecule. Potential energy curves for the A and X states were constructed up to v = 25 and v = 36 respectively, much farther than previously known. Except a drastic change in the nature of the BiF (X O <sup>+</sup> ) band strength, the dissociation energy is estimated 40 000 cm <sup>-1</sup> almost twice the calculated current value. We are currently constructing a new BiF source in order to better determine the electronic structure of low-lying states of BiF.				
14. SUBJECT TERMS Chemical laser mechanism : BiF spectra and energy levels of the A O <sup>+</sup> and X O <sup>+</sup> electronic states			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT	

AD-A233 032

91 3 15 114

EDARD TR-91-04

This report has been reviewed by EDARD and is releasable to the National Technical Information Service (NTIS). At NTIS it will be releasable to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

*Kirk E. Hackett*

KIRK E. HACKETT, Ph.D.  
Chief, Physics

*Fred T. Gilliam*

FRED T. GILLIAM, Lt Col, USAF  
Chief Scientist

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution/	
Availability	
Dist	Avail and/or Special
A-1	



**UNIVERSITE CLAUDE BERNARD - LYON I**

**LABORATOIRE DE SPECTROMETRIE  
IONIQUE ET MOLECULAIRE**

associé au C.N.R.S. U.A. 171

**EQUIPE DE SPECTROMETRIE MOLECULAIRE**

bâtiment 205

43, boulevard du 11 Novembre 1918

69622 VILLEURBANNE Cedex (France)

Tél. 72.44.80.00      Secrétariat 85.57

Télécopie (33) 78.89.44.15

Professeur R. BACIS      72.44.81.82

**Final Scientific Report - GRANT AFOSR - 88 - 0347**

During the period covered by the U.S. Air Force grant, a Broida type oven was constructed in order to investigate the spectroscopy of the low-lying states of the BiF molecule. Moderate resolution spectra of the chemiluminescence from the Bi + F<sub>2</sub> reaction were recorded by using the Fourier transform instrument by Jean Vergès at the Aimé Cotton Laboratory in Paris. These spectra allowed an extended vibrational analysis of the BiF AO<sup>+</sup> and XO<sup>+</sup> electronic states (1). From these data, potential energy curves for the A and X states were constructed up to  $v = 25$  and  $v = 36$ , respectively, much farther than previously known. LeRoy-Bernstein long range analysis applied to the vibrational constants obtained for the BiF ground state yielded an estimate for the dissociation energy of 40 000 cm<sup>-1</sup>, almost twice the current value of 21 200 cm<sup>-1</sup> (2). Since the highest observed level of the ground state,  $v = 36$ , lies only 6 400 cm<sup>-1</sup> below this energy, a drastic change in the nature of the BiF bond as it is stretched would be necessary to conform with the current value of the BiF bond strength. Only experimental observation of higher vibrational levels of the BiF X state will resolve this question.

We attempted to improve the BiF emission intensity from our Broida oven in order to extend the knowledge of the BiF ground state closer to the dissociation energy. Unfortunately, the temperature range obtainable with our current apparatus is limited to less than 600°C in the reaction zone where the major product of the Bi + F<sub>2</sub> reaction is BiF<sub>3</sub> and hence any increase in F<sub>2</sub> density results in a decrease of the BiF A-X emission intensity. However, at temperatures above 700°C, the BiF<sub>3</sub> + 2 Bi - 3 BiF reaction begins to dominate (3). We are currently constructing a new Broida oven that will operate in the 900-950°C range to exploit this reaction, so much higher BiF densities can be achieved. With this new source we will observe A-X emission to much higher X state levels. This apparatus will also operate at pressures up to several torr where several new band systems appear (4).

We have also built a BiF emission source based on microwave excitation. This apparatus consists of two separate ovens with independent temperature control in order to produce both Bi and BiF<sub>3</sub> vapor. These are entrained in a F<sub>2</sub>/He mixture and flowed through a microwave discharge cavity. The BiF A-X and B-X emission is intense while the fluorescence of the so-called C<sub>1</sub>-X<sub>3</sub> and C<sub>2</sub>-X<sub>2</sub> systems is relatively weak, as shown in Fig. 1. Assignments of the electronic states responsible for the C<sub>1</sub>-X<sub>3</sub> and C<sub>2</sub>-X<sub>2</sub> bands have been suggested in Ref. 5. We attempted to observe high resolution Fourier Transform spectra of these systems to better characterize these poorly known states of BiF. However, the presence of strong Bi resonance lines prevented observation of these weak bands under high resolution.

The X<sub>1</sub> state, predicted by ab-initio calculations (2) to lie some 7 600 cm<sup>-1</sup> above the ground state, has recently been observed by Devore et al. under low resolution conditions (4). We will attempt to obtain high resolution emission spectra using our high temperature Broida oven source and as confirm the assignments of ref. 4. The X<sub>1</sub> state may also be populated by energy transfer from the near-resonant process : BiF (XO<sup>+</sup>) + O<sub>2</sub> (<sup>1</sup>Δ<sub>g</sub>) → BiF (X<sub>1</sub>) + O<sub>2</sub> (<sup>3</sup>Σ<sub>g</sub><sup>-</sup>). We will also try to record BiF X<sub>1</sub> - XO<sup>+</sup> emission spectra in the 7600 cm<sup>-1</sup> region in order to characterize the X<sub>1</sub> state. We will also obtain IR emission spectra from the system reported in ref. 6. In addition, a King furnace is being installed to obtain BiF from the thermolysis of BiF<sub>3</sub> at temperature of 1 200-1 300° C so that high resolution Fourier transform absorption spectra can be recorded. These new investigations will yield information of the electronic structure of low-lying states of BiF as well as provide a more precise determination of the BiF bond strength.

## References

- 1 A.J. ROSS, R. BACIS, J. d'INCAN, C. EFFANTIN, B. KOFFEND, A. TOPOUZKHANIAN and J. VERGES  
Chem. Phys. Lett. 166, 539 (1990)
- 2 K. BALASUBRAMANIAN  
Chem. Phys. Lett. 127, 324 (1986)
- 3 P. KUIJPERS, A. DYMANUS  
Chem. Phys. 24, 97 (1977)
- 4 T.C. DEVORE, L. BROCK, K. DULANEY, J.L. GOLE  
Chem. Phys. Lett. to be published
- 5 A.K. CHAUDHRY, K.N. UPA DHYA, D.K. RAI  
J. Phys. B, 2, 628 (1969)
- 6 C.I. FRUM, R. ENGLEMAN, P.F. BERNATH  
J. Chem. Phys. 93, 5457 (1970)

SY0.BIFPHASE . SPC	DET: PM	SOURC: BIF	B/S: QUV	SP: 8	SM: -4	APER: 7	EV: E
RES : 61.741	DATE: 08-JUN-90	TIME: 18:35:58		BG: 1	PG: 1	APOD: Bartlett	
COADD: 32				LP: 5	HP: 2	FILTER: FI2	

BiF discharge with Moncorge's filter. PM 850V.  
Speed .2 cm/s.




Fig 1 : Fourier Transform low resolution spectrum from a microwave discharge in BiF